


MATH 223: Multivariable Calculus



Class 25: April 18, 2022




Notes on Assignment 23
Assignment 24
Jacobi's Theorem on Change of Variable

Announcements

Review Improper Integrals:


$$\int_1^{\infty} \frac{1}{x^n} dx$$



Description Due Today

Mathematician of the Week

Conel Hugh Alexander



April 19, 1909 – February 15, 1974
Cryptanalyst and Chess Champion

Biography

G. H. Hardy: "He was the only genuine mathematician I knew who did not become a professional mathematician"

This Week:
Change of Variable
Improper Integrals
Application to Probability

Change of Variable aka Method of Substitution

A common technique in the evaluation of integrals is to make a change of variable in the hopes of simplifying the problem of determining an antiderivatives



Example: Evaluate $\int_{x=0}^{x=2} \frac{2x}{1+x^2} dx$

$$\begin{array}{l} \text{Let } u = 1 + x^2 \quad \left| \quad x = 0 \rightarrow u = 1 + 0^2 = 1 \right. \\ \text{The } du = 2x dx \quad \left| \quad x = 2 \rightarrow u = 1 + 2^2 = 5 \right. \end{array}$$

$$\int_{x=0}^{x=2} \frac{2x}{1+x^2} dx = \int_{u=1}^{u=5} \frac{1}{u} du = \ln 5 - \ln 1 = \ln 5$$

$$\int_{x=0}^{x=2} \frac{2x}{1+x^2} dx = \int_{u=1}^{u=5} \frac{1}{u} du$$

Let's look at what is happening geometrically:





Not only does the function change, but also the region of integration.

The region of integration changes from an interval of length 2 to an interval of length 4.

The interval also moves to a new location.


In computing multiple integrals, the corresponding change in the region may be more complicated.

By a **change of variable**, we will mean a vector function T from \mathbb{R}^n to \mathbb{R}^n . It is convenient to use different letters to denote the spaces; e.g, $T : U^n \rightarrow \mathbb{R}^n$



Carl Gustav Jacob Jacobi

December 10, 1804 – February 18, 1851



For further information see his [Biography](#)

Jacobi's Theorem

Let \mathcal{R} be a set in \mathbb{U}^n and $T(\mathcal{R})$ its image under T ; that is,

$$T(\mathcal{R}) = \{T(\vec{u}) : \vec{u} \text{ is in } \mathcal{R}\}$$

Suppose $f : \mathbb{R}^n \rightarrow \mathbb{R}^1$ is a real-valued function.

Then, under suitable conditions,

$$\int_{T(\mathcal{R})} f(\vec{x}) dV_{\vec{x}} = \int_{\mathcal{R}} f(T(\vec{u})) |\det T'(\vec{u})| dV_{\vec{u}}$$

- ▶ T is continuous differentiable
- ▶ Boundary of \mathcal{R} is finitely many smooth curves
- ▶ T is one-to-one on interior of \mathcal{R}
- ▶ The Jacobian Determinant $\det T'$ is non zero on interior of \mathcal{R} .
- ▶ The function f is bounded and continuous on $T(\mathcal{R})$

$$\int_{T(\mathcal{R})} f(\vec{x}) dV_{\vec{x}} = \int_{\mathcal{R}} f(T(\vec{u}) | \det T'(\vec{u}) |) dV_{\vec{u}}$$

In our example: $u = 1 + x^2$ so $x = \sqrt{u-1}$

Thus $T(u) = \sqrt{u-1} = (u-1)^{1/2}$ so

$$T'(u) = \frac{1}{2}(u-1)^{-1/2} = \frac{1}{2\sqrt{u-1}}$$


$$\int_0^2 \frac{2x}{1+x^2} dx = \int_{T(\mathcal{R})} f(\vec{x}) dV_{\vec{x}} = \int_1^5 f(T(u) | \det T'(u) |) du$$

$$\text{Now } f(T(\vec{u})) = \frac{2T(u)}{1+(T(u))^2} = \frac{2\sqrt{u-1}}{1+u-1} = \frac{2\sqrt{u-1}}{u}$$

$$\det T'(u) = \left| \frac{1}{2\sqrt{u-1}} \right| = \frac{1}{2\sqrt{u-1}} \text{ so } f(T(\vec{u})) \det T'(u) = \frac{1}{u}$$

$$\text{so } \int_0^2 \frac{2x}{1+x^2} dx = \int_1^5 \frac{2\sqrt{u-1}}{u} \frac{1}{2\sqrt{u-1}} du = \int_1^5 \frac{1}{u} du$$

Example: Polar Coordinate Change of Variable

$$\mathcal{U}^2 \quad T \rightarrow \quad \mathcal{R}^2$$


$x = r \cos \theta$
 $y = r \sin \theta$

$$T(r, \theta) = (r \cos \theta, r \sin \theta)$$


$$T' = \begin{pmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix} \text{ so } \det T' = r \cos^2 \theta + r \sin^2 \theta = r$$

$$\text{Thus } \int_{T(R)} f(x, y) dx dy = \int_R f(r \cos \theta, r \sin \theta) r dr d\theta$$

$$\int_{T(R)} f(x, y) dx dy = \int_R f(r \cos \theta, r \sin \theta) r dr d\theta$$

Example: $f(x, y) = x^2 + y^2$


$$T(R) = \text{Half Disk} = \{(x, y) : -1 \leq x \leq 1, 0 \leq y \leq \sqrt{1-x^2}\}$$





$$I = \int_{-1}^1 \int_0^{\sqrt{1-x^2}} (x^2 + y^2) dy dx$$

Describe Region in Polar Coordinates: $0 \leq r \leq 1, 0 \leq \theta \leq \pi$

$$I = \int_{\theta=0}^{\pi} \int_{r=0}^1 r^2 r dr d\theta = \int_{\theta=0}^{\pi} \left. \frac{r^4}{4} \right|_0^1 d\theta = \int_{\theta=0}^{\pi} \frac{1}{4} d\theta = \frac{\pi}{4}$$



Look At This Transformation More Closely



$$\begin{aligned} A : 0 \leq r \leq 1, \theta = 0 \\ x = r \cos \theta = r \cos 0 = r \\ y = r \sin \theta = r \sin 0 = 0 \end{aligned}$$

$$\begin{aligned} B : r = 1, 0 \leq \theta \leq \pi \\ x = r \cos \theta = \cos \theta \\ y = r \sin \theta = \sin \theta \end{aligned}$$


$$\begin{aligned} C : 0 \leq r \leq 1, \theta = \pi \\ x = r \cos \theta = r \cos \pi = -r \\ y = r \sin \theta = r \sin \pi = 0 \end{aligned}$$

$$\begin{aligned} D : r = 0, 0 \leq \theta \leq \pi \\ x = r \cos \theta = 0 \\ y = r \sin \theta = 0 \end{aligned}$$

Problem: Evaluate $\iiint_C \sqrt{x^2 + y^2 + z^2} dV$

where C is the ice cream cone

$$\{(x, y, z) : x^2 + y^2 + z^2 \leq 1, x^2 + y^2 \leq \frac{z^2}{3}, z \geq 0\}$$



Example: Spherical Coordinates

$$x = r \sin \phi \cos \theta \quad T : (r, \phi, \theta) \rightarrow (x, y, z)$$

$$y = r \sin \phi \sin \theta \quad \det T' = r^2 \sin \phi$$

$$z = r \cos \phi$$

Problem: Evaluate $\iiint_C \sqrt{x^2 + y^2 + z^2} dV$

where C is the ice cream cone

$$\{(x, y, z) : x^2 + y^2 + z^2 \leq 1, x^2 + y^2 \leq \frac{z^2}{3}, z \geq 0\}$$

$$z \geq 0 \text{ implies } \phi \leq \frac{\pi}{2}$$


$$x^2 + y^2 + z^2 \leq 1 \text{ implies } r \leq 1$$

$$x^2 + y^2 \leq \frac{z^2}{3} \text{ implies } r^2 \sin^2 \phi \leq \frac{r^2 \cos^2 \phi}{3}$$

$$\text{implies } \tan^2 \phi \leq \frac{1}{3} \text{ implies } \phi \leq \frac{\pi}{6}$$

$$\int_{\theta=0}^{2\pi} \int_{\phi=0}^{\pi/6} \int_{r=0}^1 \sqrt{r^2} r^2 \sin \phi \, dr \, d\phi \, d\theta$$

Example: Evaluate $\iiint_D z^2 dV$ where D is the interior of the ellipsoid $\frac{x^2}{4} + \frac{y^2}{16} + \frac{z^2}{9} = 1$



STEP 1: Let $u = \frac{x}{2}$, $v = \frac{y}{4}$, $w = \frac{z}{3}$.

Equation of the ellipsoid becomes $u^2 + v^2 + w^2 = 1$ (unit sphere)

So $x = 2u$, $y = 4v$, $z = 3w$ gives $T(u, v, w) = (2u, 4v, 3w)$ and

$$T' = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 3 \end{pmatrix} \text{ so } \det T' = 2 \times 4 \times 3 = 24$$

Thus $\iiint_D z^2 = \iiint (3w)^2 (24) du dv dw = 216 \iiint w^2 du dv dw$

STEP 2: Switch to Spherical Coordinates:

$$u = r \sin \phi \cos \theta, v = r \sin \phi \sin \theta, w = r \cos \phi$$

$$\begin{aligned} 216 \iiint w^2 du dv dw &= 216 \iiint (r \cos \phi)^2 r^2 \sin \phi dr d\phi d\theta \\ &= 216 \int_{\theta=0}^{2\pi} \int_{\phi=0}^{\pi} \int_{r=0}^1 r^4 \cos^2 \phi \sin \phi dr d\phi d\theta \\ &= (216)(2\pi) \int_{\phi=0}^{\pi} \int_{r=0}^1 r^4 \cos^2 \phi \sin \phi dr d\phi \\ &= (216)(2\pi) \frac{1}{5} \int_{\phi=0}^{\pi} \cos^2 \phi \sin \phi d\phi \\ &= \frac{(216)(2\pi)}{5} \left[-\frac{\cos^3 \phi}{3} \right]_{\phi=0}^{\pi} = \frac{(216)(2\pi)}{5} \frac{2}{3} = \frac{288\pi}{5} \end{aligned}$$