

MATH 223: Multivariable Calculus


Divergence, Curl and Gradient Operations

(i) Divergence

➤ The divergence of a vector \mathbf{V} written as $\text{div } \mathbf{V}$ represents the

$$\text{div } \mathbf{V} = \nabla \cdot \mathbf{V} =$$

$$\frac{\partial V_x}{\partial x} + \frac{\partial V_y}{\partial y} + \frac{\partial V_z}{\partial z}$$



(a) positive divergence



(b) negative divergence



(c) zero divergence

Class 30: May 2, 2022



Notes on Assignment 28
Assignment 29
Divergence and Curl

Announcements

Exam 3: Tonight
Axinn 229 at. 7 PM
One Page of Notes

Today

Begin Chapter 8: Vector Field Theory

**Divergence and Curl: Measures of Rates of Change of
Vector Fields**

Divergence of a Vector Field

Definition $\operatorname{div} \mathbf{F} = \text{trace of } \mathbf{F}'$ of the Jacobi Matrix

Example $\mathbf{F}: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $\mathbf{F}(x, y) = (2x - y, x - 3y)$

$$\mathbf{F}' = \begin{pmatrix} 2 & -1 \\ 1 & -3 \end{pmatrix} \text{ implies } \operatorname{div} \mathbf{F} = 2 - 3 = -1$$

Example: $\mathbf{F}: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ by $\mathbf{F}(x, y, z) = (xy, yz, zx)$

$$\mathbf{F}' = \begin{pmatrix} y & - & - \\ - & z & - \\ - & - & x \end{pmatrix} \text{ implies } \operatorname{div} \mathbf{F} = y + z + x$$

Example: $\mathbf{F}: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ by $\mathbf{F}(x, y, z) = (yz, xz, xy)$

Alternate Notation: $yz\mathbf{i} + xz\mathbf{j} + xy\mathbf{k}$

$$\mathbf{F}' = \begin{pmatrix} 0 & z & y \\ z & 0 & y \\ y & z & 0 \end{pmatrix} \text{ implies } \operatorname{div} \mathbf{F} = 0$$

$$\frac{\partial}{\partial x} \quad \frac{\partial}{\partial y} \quad \frac{\partial}{\partial z}$$

$$\mathbf{F}' = \begin{matrix} F1 \\ F2 \\ F3 \end{matrix} \begin{pmatrix} 0 & z & y \\ z & 0 & y \\ y & z & 0 \end{pmatrix} \text{ implies } \operatorname{div} \mathbf{F} = 0$$

In general, **div \mathbf{F} is a real -valued function of n variables.**

Notes

1. Gauss's Theorem: $\int_R \operatorname{div} \mathbf{F} dV = \int_{\partial R} \mathbf{F} \cdot d\mathbf{S}$
2. $\operatorname{div} \mathbf{F}$ gives expansion rate of fluid at point \mathbf{x}
 $\operatorname{div} \mathbf{F} > 0$ means fluid is expanding, getting less dense
 $\operatorname{div} \mathbf{F} < 0$ means fluid is contracting, becomes more dense
3. Alternate Notation; $\mathbf{F} = (F_1, F_2, F_3)$, $\nabla = (\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z})$
Then $\operatorname{div} \mathbf{F} = \mathbf{F} \cdot \nabla$


Example

$$\mathbf{F}(x, y, z) = (xy^2 + z \ln(1 + y^2), \sin(xz) - zy, x^2z + \arctan y + e^{x^2})$$


$$\operatorname{div} \mathbf{F} = y^2 - z + x^2$$

$$\text{so } \operatorname{div} \mathbf{F} > 0 \text{ if } x^2 + y^2 > z$$

$z = x^2 + y^2$ is equation of elliptic paraboloid.




Divergence is positive on the outside, negative on the inside.



$$\mathbf{F}(x, y) = \frac{1}{8}(-x + y)\mathbf{i} + \frac{1}{8}(-x - y)\mathbf{j}$$

Area decreased: $\text{div}\mathbf{F}(x, y) = -\frac{1}{4}$

(a)



$$\mathbf{G}(x, y) = -\frac{1}{4}(y/\sqrt{x^2 + y^2})\mathbf{i} + \frac{1}{4}(x/\sqrt{x^2 + y^2})\mathbf{j}$$

Area preserved: $\text{div}\mathbf{G}(x, y) = 0$

(b)

Figure 8.10

Curl of a Vector Field

Curl measures local tendency of a vector field and its flow lines to circulate around some axis.

The curl of a vector field is itself a vector field.

Setting; $\mathbf{F} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is our vector field

$$\mathbf{F} = (F_1, F_2, F_3) \text{ so}$$

$$\mathbf{F}(x, y, z) = (F_1(x, y, z), F_2(x, y, z), F_3(x, y, z))$$

$$\text{Formal Definition: } \text{curl } \mathbf{F} = \left(\frac{\partial F_3}{\partial y} - \frac{\partial F_2}{\partial z}, \frac{\partial F_1}{\partial z} - \frac{\partial F_3}{\partial x}, \frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right)$$

Mnemonic Device:

$$\text{curl } \mathbf{F} = \det \begin{pmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_1 & F_2 & F_3 \end{pmatrix}$$

Expand along first row:

$$\text{curl } \mathbf{F} = \begin{vmatrix} \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_2 & F_3 \end{vmatrix} \mathbf{i} - \begin{vmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial z} \\ F_1 & F_3 \end{vmatrix} \mathbf{j} + \begin{vmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} \\ F_1 & F_2 \end{vmatrix} \mathbf{k}$$

$$\operatorname{curl} \mathbf{F} = \left(\frac{\partial F_3}{\partial y} - \frac{\partial F_2}{\partial z}, \frac{\partial F_1}{\partial z} - \frac{\partial F_3}{\partial x}, \frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right)$$

Example: $\mathbf{F}(x, y, z) = (xyz, y - 3z, 2y)$

$$\begin{aligned} \operatorname{curl} \mathbf{F} &= ((2y)_y - (-3z)_z, (xyz)_z - (2y)_x, (y - 3z)_x - (xyz)_y) \\ &= (2 - (-3), xy - 0, 0 - xz) = (5, xy, -xz) \end{aligned}$$

Scalar Curl for Vector Fields in Plane

$\mathbf{F} = (F, G, 0)$ where $F(x, y)$ and $G(x, y)$ are functions only of x and y .

Then $\text{curl } \mathbf{F} = (0, 0, G_x - F_y)$

Note: Curl and Conservative Vector Field


Suppose $\mathbf{F} = (F, G, 0)$ is gradient field with $\mathbf{F} = \nabla f$.

Then $F = f_x$ and $G = f_y$

In this case, $\text{Curl } \mathbf{F} = (0, 0, f_{yx} - f_{xy}) = (0, 0, 0)$
by Clairault's Theorem on Equality of Mixed Partial.

Green's Theorem in the Plane

$$\iint_D \text{curl } \mathbf{F} = \int_{\gamma} \mathbf{F}$$



D is bounded plane region.

$C = \gamma$ is piecewise smooth boundary of D

F and G are continuously differentiable functions defined on D

Then

$$\int \int (G_x - F_y) dx dy = \int_{\gamma} F dx + G y$$

where γ is parametrized so it is traced once with D on the left.

Application of Green's Theorem in the Plane

$$\iint_D \text{curl } \mathbf{F} = \int_{\gamma} \mathbf{F}$$

Example $\mathbf{F}(x, y) = (0, x)$ implies $\text{curl } \mathbf{F} = 1 - 0 = 1$

Hence $\iint_D \text{curl } \mathbf{F} = \iint_D 1 = \text{area of } D$

Green's Theorem enables us to find the area of a planar region if we can develop a parametrization of its boundary.

Example Consider the unit disk D of radius r centered at origin.

Let $g(t) = (r \cos t, r \sin t), 0 \leq t \leq 2\pi$

So $g'(t) = (-r \sin t, r \cos t)$

and $\mathbf{F}(g(t)) = (0, r \cos t)$

Then $\mathbf{F}(g(t)) \cdot g'(t) = r^2 \cos^2 t dt$

Thus area of $D = \iint_D 1 = \iint_D \text{curl } \mathbf{F} = \int_{\gamma} \mathbf{F} = \int_0^{2\pi} r^2 \cos^2 t dt$

$$\int_0^{2\pi} r^2 \cos^2 t dt = r^2 \int_0^{2\pi} \frac{1+\cos 2t}{2} dt = \frac{r^2}{2} \left[t + \frac{1}{2} \sin 2t \right]_0^{2\pi} = \pi r^2$$

Using Green's Theorem

(1) Compute $\iint_D \text{curl } \mathbf{F}$ by using $\int_\gamma \mathbf{F}$

(2) Compute $\int_\gamma \mathbf{F}$ by using $\iint_D \text{curl } \mathbf{F}$

Curl of a Vector Field

Curl measures local tendency of a vector field and its flow lines to circulate around some axis.

The curl of a vector field is itself a vector field.

Setting; $\mathbf{F} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is our vector field

$$\mathbf{F} = (F_1, F_2, F_3) \text{ so } \mathbf{F}(x, y, z) = (F_1(x, y, z), F_2(x, y, z), F_3(x, y, z))$$

$$\text{Formal Definition: } \text{curl } \mathbf{F} = \left(\frac{\partial F_3}{\partial y} - \frac{\partial F_2}{\partial z}, \frac{\partial F_1}{\partial z} - \frac{\partial F_3}{\partial x}, \frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right)$$

Mnemonic Device:

$$\text{curl } \mathbf{F} = \det \begin{pmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_1 & F_2 & F_3 \end{pmatrix}$$

Expand along first row:

$$\text{curl } \mathbf{F} = \begin{vmatrix} \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_2 & F_3 \end{vmatrix} \mathbf{i} - \begin{vmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial z} \\ F_1 & F_3 \end{vmatrix} \mathbf{j} + \begin{vmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} \\ F_1 & F_2 \end{vmatrix} \mathbf{k}$$

Scalar Curl for Vector Fields in Plane

$\mathbf{F} = (F, G, 0)$ where $F(x, y)$ and $G(x, y)$ are functions only of x and y .

$$\text{Then curl } \mathbf{F} = (0, 0, G_x - F_y)$$

Note: Curl and Conservative Vector Field


Suppose $\mathbf{F} = (F, G, 0)$ is gradient field with $\mathbf{F} = \nabla f$.

$$\text{Then } F = f_x \text{ and } G = f_y$$

In this case, $\text{Curl } \mathbf{F} = (0, 0, f_{yx} - f_{xy}) = (0, 0, 0)$
by Clairaut's Theorem on Equality of Mixed Partial.

Green's Theorem in the Plane

$$\iint_D \text{curl } \mathbf{F} = \int_{\gamma} \mathbf{F}$$



D is bounded plane region.

$C = \gamma$ is piecewise smooth boundary of D

F and G are continuously differentiable functions defined on D

Then

$$\int \int (G_x - F_y) dx dy = \int_{\gamma} F dx + G y$$

where γ is parametrized so it is traced once with D on the left.

Using Green's Theorem

(1) Compute $\iint_D \text{curl } \mathbf{F}$ by using $\int_\gamma \mathbf{F}$


(2) Compute $\int_\gamma \mathbf{F}$ by using $\iint_D \text{curl } \mathbf{F}$

Example

Find

$$\int_{\gamma} (1+10xy+y^2)dx + (6xy+5x^2)dy = \int_{\gamma} (1+10xy+y^2, 6xy+5x^2)$$

where γ is boundary of the rectangle with vertices $(0,0)$, $(2,0)$, $(2,1)$, and $(0,1)$.



Note: Direct Computation requires 4 integrals.

$$F(x, y) = 1 + 10xy + y^2. \quad G(x, y) = 6xy + 5x^2$$

$$F_y = 10x + 2y \quad . \quad G_x = 6y + 10x$$

$$G_x - F_y = 6y + 10x - 10x - 2y = 4y$$

$$\int_{\gamma} \mathbf{F} = \iint_D \text{curl } \mathbf{F} = \int_0^2 \int_0^1 4y \, dy \, dx = \int_0^2 [2y^2]_0^1 = \int_0^2 2 \, dx = 4$$



George Green
1793 – 1841



Mikhail Ostrogradsky
1801 – 1861

