

MATH 226 Differential Equations



Class 33: May 9, 2022



Notes on Assignment 21
Assignment 22
Sample Final Exam

Announcements

Project 3 Due Friday

Course Response Forms
In Class Next Monday
Bring Laptop/SmartPhone

**Final Exam
Friday, May 20: 9 - Noon**

Numerical Approximations To Solutions of Differential Equations

$$y' = f(t, y) \text{ with } y(t_0) = y_0$$

Euler's Method: $y_{n+1} = y_n + f(t_n, y_n)$

Improved Euler's:

$$y_{n+1} = y_n + \frac{h}{2} [f(t_n, y_n) + f(t_n + h, y_n + f(t_n, y_n))]$$

Second Order Methods

$$y_{n+1} = y_n + h(ak_1 + bk_2) \text{ where}$$

$$k_1 = f(t_n, y_n)$$

$$k_2 = f(t_n + ph, y_n + qk_1 h)$$

$$\text{with } a + b = 1, bp = bq = \frac{1}{2}$$

Heun	$a = b = 1/2$	$p = q = 1$
Midpoint	$a = 0, b = 1$	$p = q = 1/2$
Ralston	$a = 1/4, b = 3/4$	$p = q = 2/3$

Start with the Differential Equation


Solution ϕ has $\phi'(t) = f(t_n, \phi(t))$

Integrate both sides over the interval $[t_n, t_{n+1}]$:


$$\int_{t_n}^{t_{n+1}} \phi'(t) dt = \int_{t_n}^{t_{n+1}} f(t_n, \phi(t)) dt$$

Now left hand side is $\phi(t_{n+1}) - \phi(t_n)$


and right hand side is area under curve



Thus $\phi(t_{n+1}) - \phi(t_n) = \text{Area Under Curve}$




Approximate by Area of Green Rectangle $(f(t_n, \phi(t_n))(t_{n+1} - t_n)$



$$\phi(t_{n+1}) - \phi(t_n) = \text{Area Under Curve}$$


A Better Way to Estimate Area Under the Curve




$$\text{Trapezoid Rule: } A = \frac{1}{2} [f(t_n, \phi(t_n)) + f(t_{n+1}, \phi(t_{n+1}))] (t_{n+1} - t_n)$$

$$\text{Thus } \phi(t_{n+1}) - \phi(t_n) \approx \frac{1}{2} [f(t_n, \phi(t_n)) + f(t_{n+1}, \phi(t_{n+1}))] (t_{n+1} - t_n)$$

Other Ways To Estimate Area Under Curve



Midpoint Rule
 $(b - a)f(m)$



Trapezoid Rule
 $\frac{b-a}{2} [f(a) + f(b)]$

Midpoint Rule: $(b - a)f(m)$

Trapezoidal Rule: $\frac{b - a}{2} [f(a) + f(b)]$

Weighted Average of Midpoint Rule and Trapezoidal Rule

Give Weight 2/3 to Midpoint and 1/3 to Trapezoid

$$(b - a) \left[\frac{2}{3}f(m) + \frac{1}{3} \frac{f(a) + f(b)}{2} \right] = \frac{b - a}{3} \left[\frac{4f(m) + f(a) + f(b)}{2} \right]$$

$$\frac{b - a}{6} [f(a) + 4f(m) + f(b)] = \frac{h}{3} [f(a) + 4f(m) + f(b)]$$

$$h = \frac{b - a}{2} \text{ and } m = \frac{a + b}{2}$$



Thomas Simpson
August 20, 1710 – May 14, 1761
Simpson's Rule



Johannes Kepler
Dec. 27, 1571 – Nov. 15, 1630
Keplersche Fassregel

Approximating f on $[a, b]$ with parabola

Given $f(a) = L, f\left(\frac{a+b}{2}\right) M, f(b) = R$

With $y = Ax^2 + Bx + C$

$$a^2A + aB + 1C = L$$

$$\left(\frac{a+b}{2}\right)^2 A + \left(\frac{a+b}{2}\right) B + 1C = M$$


$$b^2A + bB + 1C = R$$

$$\begin{pmatrix} a^2 & a & 1 \\ \left(\frac{a+b}{2}\right)^2 & \left(\frac{a+b}{2}\right) & 1 \\ b^2 & b & 1 \end{pmatrix} \begin{pmatrix} A \\ B \\ C \end{pmatrix} = \begin{pmatrix} L \\ M \\ R \end{pmatrix}$$

With $a = 0$ and $b = 1$, we get

$$A = 2L - 4M + 2R \quad B = -3L - R + 4M \quad C = L$$

Example: Parabola Approximation of $\sin x$ on $[0, \pi]$



$$\begin{aligned}\sin x \\ -\frac{4}{\pi^2}x^2 + \frac{4}{\pi}x\end{aligned}$$

Approximate function by the quadratic polynomial (i.e. parabola) $P(x)$ that takes the same values as the function at the end points a and b and the midpoint $m = (a + b)/2$.

We obtain $P(x) =$

$$f(a) \frac{(x - m)(x - b)}{(a - m)(a - b)} + f(m) \frac{(x - a)(x - b)}{(m - a)(m - b)} + f(b) \frac{(x - m)(x - a)}{(b - a)(b - m)}$$

Then $\int_a^b P(x) = \frac{b - a}{6} [f(a) + 4f(m) + f(b)]$

The Classic Runge–Kutta Method (RK4)

$$y_{n+1} = y_n + \frac{1}{6} (k_1 + 2k_2 + 2k_3 + k_4)$$


where

$$k_1 = f(t_n, y_n)$$


$$k_2 = f\left(t_n + \frac{h}{2}, y_n + \frac{k_1}{2}\right)$$

$$k_3 = f\left(t_n + \frac{h}{2}, y_n + h\frac{k_2}{2}\right)$$

$$k_4 = f(t_n + h, y_n + hk_3)$$



Motivation: Simpson's Rule



$$\int_a^b f(x) dx \approx \frac{h}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right]$$

More General Runge–Kutta Methods

$$y_{n+1} = y_n + h \sum_{i=1}^s b_i k_i,$$

where

$$k_1 = f(t_n, y_n),$$

$$k_2 = f(t_n + c_2 h, y_n + h(a_{21} k_1)),$$

$$k_3 = f(t_n + c_3 h, y_n + h(a_{31} k_1 + a_{32} k_2)),$$

⋮

$$k_s = f(t_n + c_s h, y_n + h(a_{s1} k_1 + a_{s2} k_2 + \cdots + a_{s,s-1} k_{s-1})).$$



Carl David Tolm  Runge
August 30, 1856
– January 3, 1927
[Runge Biography](#)



Martin Wilhelm Kutta
November 3, 1867
– December 25, 1944
[Kutta Biography](#)

Error Estimates Proportional to Step Size h

Method	Local Error	Global Error
Euler	h^2	h
Improved Euler	h^3	h^2
Runge–Kutta	h^5	h^4

Example: $y' = 1 - t + 4y$ with $y(0) = 1$

Predict Value of y at $t = 2$

Improved Euler	Runge–Kutta	Runge–Kutta	Runge–Kutta	Exact
$h = .0025$	$h = 0.2$	$h = 0.1$	$h = .05$	
$h = 1/40$	$h = 1/5$	$h = 1/10$	$h = 1/20$	
3496.6702	3490.5574	3535.8667	3539.8804	3540.2

Comparing Improved Euler and Runge–Kutta 160 Functional Evaluations

Improved Euler ($h = .025$) 1.23% Error

Runge–Kutta ($h = .05$) .00903% Error

Method	h	Evaluations	Percent Error
Improved Euler	.025	160	1.23%
Runge-Kutta	.2	40	1.4%

Runge–Kutta produces
Better Results With Similar Effort
Similar Results with Less Effort

Taylor Series Approach

$$y(t) = y(t_n) + y'(t_n)(t - t_n) + \frac{y''(t_n)}{2!}(t - t_n)^2 + \frac{y^{(3)}(t_n)}{3!}(t - t_n)^3 + \dots$$