


MATH 226: Differential Equations



February 25, 2022



Notes on Assignment 3

Assignment 4

Maple Examples (on line)

Intro to MATLAB

Announcements

First Team Project

**Examination of Models of form $P' = rP^k$ with applications to
United States Population Growth
Coronavirus Virus Deaths**

Submit Teammate Preferences by next Tuesday
Teams Assigned next Wednesday, March 2
Projects Due Friday, March 11

Today Part I

Variables Separable

$$y'(t) = f(y)g(t)$$

- ▶ Separate Variables
- ▶ Integrate
- ▶ Use Initial Value To Find Integration Constant C
- ▶ Solve for y
- ▶ Determine Interval of Validity of Solution

Today Part II

First Order LINEAR Differential Equations

$$\frac{dy}{dt} + p(t)y = g(t)$$

Method of Integrating Factors

First Order LINEAR Differential Equations

$$\frac{dy}{dt} + p(t)y = g(t)$$

$$y' + p(t)y = g(t)$$

- ▶ y' and y appear all by themselves
- ▶ No terms like y^2 or $\cos y$ or $y y'$
- ▶ $p(t)$ and $g(t)$ can be nonlinear, complicated, but continuous.

First Order LINEAR Differential Equations

$$y' + p(t)y = g(t)$$

Solution By Method of Integrating Factors:

**Multiply Equation By Factor That Converts
Left Hand Side Into a Derivative**

Fundamental Theorem of Calculus

$$\text{If } f(t) = \int p(t)dt, \text{ then } f'(t) = p(t)$$

$$\text{If } f(t) = \int_0^t p(s)ds, \text{ then } f'(t) = p(t)$$

Application:

Find the derivative of $e^{\int p(t) dt} = \exp(\int p(t) dt)$ with respect to t

Solution : Use Product Rule

$$\left(e^{\int p(t) dt} \right)' = e^{\int p(t) dt} p(t) = p(t) e^{\int p(t) dt}$$

Let's Do a Specific Example (Problem 15 in Text):

$$\text{Solve } ty' + 4y = t^2 - t + 1, \text{ with } y(1) = \frac{1}{4}$$

$$\text{Put in Standard Form (*): } y' + \frac{4}{t}y = t - 1 + \frac{1}{t}$$

$$\text{Integrating Factor is } e^{\int \frac{4}{t} dt} = e^{4 \ln t} = e^{\ln t^4} = t^4$$

$$\text{Multiply (*) by } t^4 : t^4 y' + 4t^3 y = t^5 - t^4 + t^3$$

$$(t^4 y)' = t^5 - t^4 + t^3 \text{ and integrate}$$

$$t^4 y = \frac{t^6}{6} - \frac{t^5}{5} + \frac{t^4}{4} + C$$

$$\text{Solve for } y : y = \frac{t^2}{6} - \frac{t}{5} + \frac{1}{4} + Ct^{-4}$$

Original Initial Value Problem

Solve $ty' + 4y = t^2 - t + 1$, with $y(1) = \frac{1}{4}$

Solution

$$y = \frac{t^2}{6} - \frac{t}{5} + \frac{1}{4} + \frac{C}{t^4}$$

Use initial condition to find C:

$$\frac{1}{4} = \frac{1}{6} - \frac{1}{5} + \frac{1}{4} + C$$

$$C = \frac{1}{30}$$

so

$$y = \frac{t^2}{6} - \frac{t}{5} + \frac{1}{4} + \frac{1}{30t^4}$$

General Case:

$$y' + p(t)y = g(t)$$

Multiply through by $e^{\int p(t) dt}$:

$$e^{\int p(t) dt} y' + p(t)e^{\int p(t) dt} y = g(t)e^{\int p(t) dt}$$

Rewrite Left Hand Side:

$$\left(e^{\int p(t) dt} y \right)' = g(t)e^{\int p(t) dt}$$

Integrate Both Sides:

$$e^{\int p(t) dt} y = \int g(t)e^{\int p(t) dt} + C$$

Divide by Coefficient of y :

$$y = e^{-\int p(t) dt} \int g(t)e^{\int p(t) dt} + Ce^{-\int p(t) dt}$$

$y' + p(t)y = g(t)$ has solution

$$y = e^{-\int p(t) dt} \int g(t) e^{\int p(t) dt} + C e^{-\int p(t) dt}$$



Problem 35: Construct a first order linear differential equation whose solutions are asymptotic to the line $y = 4 - t$ as $t \rightarrow \infty$.

Solution: Add a term that goes to 0 as $t \rightarrow \infty$.

One choice would be Ce^{-t} for an arbitrary constant C .

Then solution has form $y = Ce^{-t} + 4 - t$.

Differentiate with respect to t : $y' = -Ce^{-t} - 1$

But $Ce^{-t} = y - 4 + t$

So $y' = -(y - 4 + t) - 1 = -y + 4 - t - 1 = -y + 3 - t$

$$y' + y = 3 - t$$

Next Time

Modeling With First Order Differential Equations

